

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/535	A1	(11) International Publication Number: WO 99/64008 (43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/GB99/01808 (22) International Filing Date: 8 June 1999 (08.06.99) (30) Priority Data: 9812664.2 11 June 1998 (11.06.98) GB (71) Applicant (for all designated States except US): MERCK SHARP & DOHME LIMITED [GB/GB]; Hertford Road, Hoddesdon, Hertfordshire EN11 9BU (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): RUPNIAK, Nadia, Melanie [GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB). (74) Agent: HISCOCK, Ian, James; Merck & Co., Inc., European Patent Dept., Terlings Park, Eastwick Road, Harlow, Essex CM20 2QR (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: USE OF A NK-1 RECEPTOR ANTAGONIST FOR TREATING SEXUAL DYSFUNCTIONS (57) Abstract The present invention provides the use of a NK-1 receptor antagonist for the manufacture of a medicament adapted for oral administration for the treatment or prevention of sexual dysfunctions, methods of treatment using such a NK-1 receptor antagonist and pharmaceutical compositions comprising the same.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF A NK-1 RECEPTOR ANTAGONIST
FOR TREATING SEXUAL DYSFUNCTIONS

This invention relates to the treatment or prevention of sexual
5 dysfunctions by the administration of a NK-1 receptor antagonist, in
particular, 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-
3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a
pharmaceutically acceptable salt thereof.

Sexual dysfunction is characterised by a disturbance in the
10 processes that are involved in the sexual response cycle or by pain
associated with sexual intercourse. The sexual response cycle comprises
the four phases of desire, excitement, orgasm and resolution. Disorders of
sexual response may occur at one or more of these phases.

Sexual dysfunctions cause marked distress and interpersonal
15 difficulty. The sexual dysfunctions include sexual desire disorders, sexual
arousal disorders, orgasmic disorders and sexual pain disorders. None of
these disorders is adequately treated with existing therapies. Thus, for
example, male erectile disorder (a sexual arousal disorder) is currently
treated using vasoactive drugs such as papaverine or prostaglandin E1
20 which are injected directly into the corpus cavernosum. This treatment is
undesirable, both due to the undesirable side-effects associated with the
therapeutic agents and due to the methodology employed. Where oral
therapeutic agents have been tested, these have generally proved
unsatisfactory because of their poor efficacy, their safety, or undesirable
25 side-effects.

Neurokinin 1 (NK-1; substance P) receptor antagonists are being
developed for the treatment of a number of physiological disorders
associated with an excess or imbalance of tachykinins, and in particular
substance P. Examples of conditions in which substance P has been
30 implicated include disorders of the central nervous system such as

anxiety, depression and psychosis (see, for instance, International (PCT) patent specification Nos. WO 95/16679, WO 95/18124 and WO 95/23798).

More recently, International (PCT) patent specification No. WO 96/24353 (published 15th August 1996) suggests that a more efficacious
5 and safe treatment of psychiatric disorders would be achieved using a combination of a tachykinin antagonist and a serotonin agonist or selective serotonin reuptake inhibitor (SSRI). However, such as regimen would not be free of side-effects due to the serotonin agonist or SSRI.

In view of the short-comings of existing therapy, there is a need for
10 new, safe and effective treatment for sexual dysfunctions.

The present invention provides the use of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof in an oral, once-a-day medicament for the treatment of sexual
15 dysfunctions. The compounds of this class advantageously exhibit a rapid onset of action and a favourable side-effect profile.

The exceptional pharmacology of the NK-1 receptor antagonist of use in the present invention enables the treatment of sexual dysfunctions, without the need for concomitant therapy, and in particular, without the
20 need for concomitant use of a serotonin agonist or an SSRI.

Furthermore, the exceptional pharmacology of the NK-1 receptor antagonist of use in the present invention results in a rapid onset of action.

The present invention accordingly provides the use of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-
25 4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament adapted for oral administration for the treatment or prevention of sexual dysfunctions.

The present invention also provides a method for the treatment or
30 prevention of sexual dysfunctions, which method comprises the oral administration to a patient in need of such treatment of an effective

amount of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof.

In a further aspect of the present invention, there is provided an oral pharmaceutical composition for the treatment of sexual dysfunctions which comprises 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier or excipient.

As used herein, the term "sexual dysfunctions" includes sexual desire disorders, sexual arousal disorders, orgasmic disorders, sexual pain disorders, sexual dysfunction due to a general medical condition, substance-induced sexual dysfunction and sexual dysfunction not otherwise specified. These sexual dysfunctions may be further defined by the nature of the onset of the disorder: either lifelong type or acquired type; by the context in which the disorder occurs: either generalized type or situational type; and by the etiological factors associated with the disorder: either due to psychological factors or due to combined factors.

Specifically, sexual desire disorders include hypoactive sexual desire disorder and sexual aversion disorder. Sexual arousal disorders include female sexual arousal disorder and male erectile disorder. Orgasmic disorders include female orgasmic disorder, male orgasmic disorder and premature ejaculation. Sexual pain disorders include dyspareunia and vaginismus. Sexual dysfunctions due to a general medical condition may result from neurological conditions (e.g. multiple sclerosis, spinal cord lesions, neuropathy and temporal lobe lesions), endocrine conditions (e.g. diabetes melitus, hypothyroidism, hypogonadal states and pituitary dysfunction), and vascular conditions and genitourinary conditions (e.g. testicular disease, Peyronie's disease, urethral infections, postprostatectomy complications, genital injury or infection, atrophic vaginitis, infections of the vagina and external

genitalia, postsurgical complications such as episiotomy scars, shortened vagina, cystitis, endometriosis, uterine prolapse, pelvic infections and neoplasms). Substance-induced sexual dysfunction can occur in association with intoxication with the following classes of substance:

5 alcohol; amphetamine (and amphetamine-like substances); cocaine; opioids; sedatives, hypnotic and anxiolytics; and other unknown substances. A decrease in sexual interest and orgasmic disorders may also be caused by prescribed medication including antihypertensives, histamine H₂-receptor antagonists, antidepressants, neuroleptics,

10 anxiolytics, anabolic steroids, and antiepileptics. Painful orgasm has been reported with fluphenazine, thioridazine and amoxapine. Priapism has been reported with the use of chlorpromazine, trazodone and clozapine, and following penile injections of papaverine or prostaglandin. Selective serotonin reuptake inhibitors may cause decreased sexual desire or

15 arousal disorders.

Also, as used herein, the term "sexual dysfunctions" includes any of the aforementioned sexual dysfunctions, including loss of libido, resulting from other medical conditions, most especially resulting from depression and/or anxiety.

20 As used herein, the term "treatment" refers both to the treatment and to the prevention or prophylactic therapy of the aforementioned conditions.

In particular, the NK-1 receptor antagonist of use in the present invention is the compound 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-

25 hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof.

Full descriptions of the preparation of the NK-1 receptor antagonist which may be employed in the present invention may be found in International Patent Specification No. WO 95/18124 and US Patent No.

30 5,612,337.

Suitable pharmaceutically acceptable salts of the NK-1 receptor antagonist of use in the present invention include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable non-toxic acid such as
5 hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid or sulphuric acid. Salts of amine groups may also comprise the quaternary ammonium salts in which the amino nitrogen atom carries an alkyl, alkenyl, alkynyl or aralkyl group.

10 Preferably the compositions containing the NK-1 receptor antagonist of use according to the present invention are in unit dosage forms such as tablets, pills, capsules, wafers and the like. Additionally, the NK-1 receptor antagonist of use according to the present invention may be presented as granules or powders for extemporaneous formulation
15 as volume defined solutions or suspensions. Alternatively, the NK-1 receptor antagonist of use according to the present invention may be presented in ready-prepared volume defined solutions or suspensions. Preferred forms are tablets and capsules.

For preparing solid compositions such as tablets, the principal
20 active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the
25 present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules.
30 This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of

the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, peanut oil or soybean oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

Compositions of the present invention may also be administered via the buccal cavity using conventional technology, for example, absorption wafers.

Compositions in the form of tablets, pills, capsules or wafers for oral administration are particularly preferred.

A minimum dosage level for the NK-1 receptor antagonist is about 1mg per day, preferably about 5mg per day and especially about 10mg per day. A maximum dosage level for the NK-1 receptor antagonist is about 1500mg per day, preferably about 1000mg per day and especially about 500mg per day. The compounds are administered once a day.

It will be appreciated that the amount of the NK-1 receptor antagonist required for use in the treatment or prevention of sexual dysfunctions will vary not only with the particular compounds or compositions selected but also with the route of administration, the nature of the condition being treated, and the age and condition of the patient, and will ultimately be at the discretion of the patient's physician or pharmacist.

The activity of the NK-1 receptor antagonist of use in the present invention may be measured using the following assays:

10

ASSAY 1: NK-1 Receptor binding

NK-1 receptor binding assays are performed in intact Chinese hamster ovary (CHO) cells expressing the human NK-1 receptor using a modification of the assay conditions described by Cascieri *et al*, *J. Pharmacol. Exp. Ther.*, 1992, 42, 458. Typically, the receptor is expressed at a level of 3×10^5 receptors per cell. Cells are grown in monolayer culture, detached from the plate with enzyme-free dissociation solution (Speciality Media Inc.), and washed prior to use in the assay. ^{125}I -Tyr⁸-substance P (0.1nM, 2000Ci/mmol; New England Nuclear) is incubated in the presence or absence of test compounds (dissolved in 5 μl dimethylsulphoxide, DMSO) with 5×10^4 CHO cells. Ligand binding is performed in 0.25ml of 50mM Tris-HCl, pH7.5, containing 5mM MnCl_2 , 150mM NaCl, 0.02% bovine serum albumin (Sigma), 50 $\mu\text{g/ml}$ chymostatin (Peninsula), 0.1nM phenylmethylsulphonyl fluoride, 2 $\mu\text{g/ml}$ pepstatin, 2 $\mu\text{g/ml}$ leupeptin and 2.8 $\mu\text{g/ml}$ furoyl saccharine. The incubation proceeds at room temperature until equilibrium is achieved (>40 minutes) and the receptor-ligand complex is harvested by filtration over GF/C filters pre-soaked in 0.1% polyethylenimine using a Tomtek 96-well harvester. Non-specific binding is determined using excess substance P (1 μM) and represents <10% of total binding.

30

ASSAY 2: Gerbil Foot-Tapping

CNS-penetrant NK-1 receptor antagonists for use in the present invention can be identified by their ability to inhibit foot tapping in gerbils induced by anxiogenic agents (such as pentagastrin) or central infusion of
5 NK-1 receptor agonists such as GR73632, or caused by aversive stimulation such as foot shock or single housing, based on the method of Rupniak & Williams, *Eur. J. Pharmacol.*, 1994, 265, 179.

Male or female Mongolian gerbils (35-70g) are anaesthetised by inhalation of an isoflurane/oxygen mixture to permit exposure of the
10 jugular vein in order to permit administration of test compounds or vehicle in an injection volume of 5ml/kg i.v. Alternatively, test compounds may be administered orally or by subcutaneous or intraperitoneal routes. A skin incision is then made in the midline of the scalp to expose the skull. An anxiogenic agent (e.g. pentagastrin) or a selective NK-1 receptor agonist
15 (e.g. GR73632 (d Ala[L-Pro⁹,Me-Leu¹⁰]-substance P-(7-11)) is infused directly into the cerebral ventricles (e.g. 3pmol in 5µl i.c.v., depending on test substance) by vertical insertion of a cuffed 27 gauge needle to a depth of 4.5mm below bregma. The scalp incision is closed and the animal
20 allowed to recover from anaesthesia in a clear perspex observation box (25cm x 20cm x 20cm). The duration and/or intensity of hind foot tapping is then recorded continuously for approximately 5 minutes. Alternatively, the ability of test compounds to inhibit foot tapping evoked by aversive stimulation, such as foot shock or single housing, may be studied using a
25 similar method of quantification.

ASSAY 3: Ferret Emesis

Individually housed male ferrets (1.0 -2.5 kg) are dosed orally by gavage with test compound. Ten minutes later they are fed with approximately 100g of tinned cat food. At 60 minutes following oral
30 dosing, cisplatin (10mg/kg) is given i.v. *via* a jugular vein catheter inserted under a brief period of halothane anaesthesia. The catheter is

then removed, the jugular vein ligated and the skin incision closed. The ferrets recover rapidly from the anaesthetic and are mobile within 10-20 minutes. The animals are observed continuously during recovery from the anaesthetic and for 4 hours following the cisplatin injection, after which
5 time the animals are killed humanely. The numbers of retches and vomits occurring during the 4 hours after cisplatin administration are recorded by trained observers.

ASSAY 4: Separation-Induced Vocalisation

10 Male and female guinea-pigs pups are housed in family groups with their mothers and littermates throughout the study. Experiments are commenced after weaning when the pups are 2 weeks old. Before entering an experiment, the pups are screened to ensure that a vigorous
15 vocalisation response is reproducibly elicited following maternal separation. The pups are placed individually in an observation cage (55cm x 39cm x 19cm) in a room physically isolated from the home cage for 15 minutes and the duration of vocalisation during this baseline period is recorded. Only animals which vocalise for longer than 5 minutes are
20 employed for drug challenge studies (approximately 50% of available pups may fail to reach this criterion). On test days each pup receives an oral dose or an s.c. or i.p. injection of test compound or vehicle and is then immediately returned to the home cage with its mother and siblings for 30 to 60 minutes (or for up to 4 hours following an oral dose, dependent upon the oral pharmacokinetics of the test compound) before social isolation for
25 15 minutes as described above. The duration of vocalisation on drug treatment days is expressed as a percentage of the pre-treatment baseline value for each animal. The same subjects are retested once weekly for up to 6 weeks. Between 6 and 8 animals receive each test compound at each dose tested.

As used herein, the term "CNS-penetrant" refers to NK-1 receptor antagonists which are able to inhibit NK-1 receptor antagonist-induced foot-tapping in the gerbil as hereinafter defined.

Essentially, hind foot-tapping in the gerbil induced by infusion of the NK-1 receptor agonist, GR73632 (d Ala[L-Pro⁹,Me-Leu¹⁰]-substance P- (7-11)), under anaesthesia, directly into the central ventricles is inhibited when a CNS-penetrant NK-1 receptor antagonist is administered intravenously immediately prior to GR73632 challenge, wherein hind foot-tapping over a period of five minutes following recovery from the anaesthesia is inhibited with an ID₅₀≤3mg/kg, and preferably with an ID₅₀≤1mg/kg.

In an alternative method, the NK-1 receptor antagonist is administered orally, 1 hour prior to GR73632 challenge, wherein the foot-tapping over a period of five minutes following recovery from anaesthesia is inhibited with an ID₅₀≤30mg/kg, and preferably with an ID₅₀≤10mg/kg.

CNS-penetrant NK-1 receptor antagonists of use in the present invention are also effective in the attenuation of separation-induced vocalisations by guinea-pig pups as hereinafter defined.

Essentially, a vocalisation response in guinea-pig pups is induced by isolation from their mothers and littermates, which response is attenuated when a CNS-penetrant NK-1 receptor antagonist is administered subcutaneously 30 minutes prior to isolation, wherein vocalisations during the first 15 minutes of isolation are attenuated with an ID₅₀≤20mg/kg, preferably with an ID₅₀≤10mg/kg, and especially with an ID₅₀≤5mg/kg.

In an alternative method, the NK-1 receptor antagonist is administered orally, 4 hours prior to isolation, wherein vocalisations during the first 15 minutes of isolation are attenuated with an ID₅₀≤20mg/kg, preferably with an ID₅₀≤10mg/kg, and especially with an ID₅₀≤5mg/kg.

A suitable selection cascade for NK₁ antagonists of use according to the present invention is as follows:

(i) Determine affinity for human NK₁ receptor in radioligand binding studies (Assay 1); select compounds with IC₅₀ ≤ 10nM, preferably
5 IC₅₀ ≤ 2nM, especially IC₅₀ ≤ 1nM.

(ii) Determine ability of compounds to penetrate CNS by their ability to inhibit foot tapping in gerbils induced by central injection of an NK₁ agonist (Assay 2); select compounds that inhibit foot tapping with
10 ID₅₀ ≤ 3mg/kg i.v., and preferably ID₅₀ ≤ 1mg/kg i.v. when administered immediately prior to central NK₁ agonist challenge, or ID₅₀ ≤ 30mg/kg p.o., and preferably ID₅₀ ≤ 10mg/kg p.o. 1 hour prior to challenge.

(iii) Determine central duration of action of compounds in gerbil foot tapping assay following intravenous administration 24 hours prior to central NK₁ agonist challenge; select compounds showing ≤ 25-fold loss of
15 potency compared with ID₅₀ determined in step (ii) above with the proviso that ID₅₀ ≤ 10mg/kg i.v., and preferably ≤ 5mg/kg i.v. after 24 hour pre-treatment.

(iv) Determine oral bioavailability of compounds by pharmacokinetic analysis, activity in gerbil foot tapping assay following
20 oral administration and/or by ability to inhibit cisplatin-induced emesis in ferrets (Assay 3); select compounds with ID₉₀ ≤ 3mg/kg p.o., and preferably ID₉₀ ≤ 1mg/kg p.o.

Particularly preferred compounds of use in the present invention are identified using steps (i) to (iv) followed by step (v):

25 (v) Determine activity of compounds in assays sensitive to conventional antidepressant/anxiolytic drugs (inhibition of pharmacologically evoked foot tapping in gerbils and/or inhibition of distress vocalisations in guinea-pig pups (Assay 4)). Select compounds with ID₅₀ ≤ 20mg/kg, and preferably ID₅₀ ≤ 10mg/kg.

30 Yet further preferred compounds of use in the present invention may be selected from those compounds which satisfy the NK-1

receptor binding criteria of step (i) which, in addition, have ≤ 5 -fold shift in affinity when incubated in the presence of human serum albumin (HSA) to show non-specific protein binding.

The NK-1 receptor antagonist of use in the present invention is the
 5 compound 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-
 3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, the
 preparation of which is described in International Patent Specification No.
 WO 95/18124 and US Patent No. 5,612,337. In the aforementioned
 assays, this compound has the following activity:

10

human NK-1 receptor binding:	IC ₅₀ = 0.12 nM
gerbil foot-tapping (5 mins.):	ID ₅₀ = 0.38 mg/kg i.v.
gerbil foot-tapping (24 hrs.):	ID ₅₀ = 2.2 mg/kg i.v.
ferret emesis:	ID ₉₀ = 1 mg/kg p.o.
guinea-pig vocalisation	
(4 hr. pre-treatment):	ID ₅₀ = 0.91 mg/kg p.o.

The following example illustrates pharmaceutical compositions according to the invention.

15 EXAMPLE 1 Tablets containing 50-300mg of NK-1 antagonist

	<u>Amount mg</u>		
NK-1 antagonist	50.0	100.0	300.0
Microcrystalline cellulose	80.0	80.0	80.0
Modified food corn starch	80.0	80.0	80.0
Lactose	189.5	139.5	139.5
Magnesium Stearate	0.5	0.5	0.5

The active ingredient, cellulose, lactose and a portion of the corn starch are mixed and granulated with 10% corn starch paste. The resulting granulation is sieved, dried and blended with the remainder of

the corn starch and the magnesium stearate. The resulting granulation is then compressed into tablets containing 50mg, 100mg and 300mg of the NK-1 receptor antagonist per tablet.

CLAIMS

1. Use of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament adapted for oral administration for the treatment or prevention of sexual dysfunctions.
2. An oral pharmaceutical composition for the treatment of sexual dysfunctions which comprises 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier or excipient.
3. A method for the treatment or prevention of sexual dysfunctions, which method comprises the oral administration to a patient in need of such treatment of an effective amount of 2-(R)-(1-(S)-(3,5-bis(trifluoromethyl)phenyl)-2-hydroxyethoxy)-3-(S)-(4-fluorophenyl)-4-(1,2,4-triazol-3-yl)methylmorpholine, or a pharmaceutically acceptable salt thereof.
4. A use according to claim 1, or a composition according to claim 2 or a method according to claim 3 wherein the sexual dysfunctions are chosen from sexual desire disorders, sexual arousal disorders, orgasmic disorders, sexual pain disorders, sexual dysfunctions due to a general medical condition, substance-induced sexual dysfunction and sexual dysfunction not otherwise specified.

INTERNATIONAL SEARCH REPORT

International Application No.

PLT/GB 99/01808

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/535

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 98 24442 A (MERCK SHARP & DOHME LIMITED) 11 June 1998 (1998-06-11) claims 1-4,6 ---	1-4
Y	WO 95 18124 A (MERCK SHARP & DOHME) 6 July 1995 (1995-07-06) cited in the application claims 9,16-18,22 page 19, line 12 - line 13 ---	1-4
Y	WO 97 18206 A (MERCK SHARP & DOHME LIMITED) 22 May 1997 (1997-05-22) claims 1-15,20 page 16, line 15 -----	1-4

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

13 September 1999

Date of mailing of the international search report

21/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 99/01808

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 3
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 99/01808

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9824442 A	11-06-1998	AU 3102097 A	14-01-1998
		AU 5485898 A	29-06-1998
		AU 5558998 A	29-06-1998
		AU 5559098 A	29-06-1998
		AU 5559198 A	29-06-1998
		AU 5559298 A	29-06-1998
		AU 5559398 A	29-06-1998
		AU 5559498 A	29-06-1998
		AU 5561398 A	29-06-1998
		AU 5752798 A	29-06-1998
		AU 5753098 A	29-06-1998
		CZ 9804247 A	12-05-1999
		WO 9824438 A	11-06-1998
		WO 9824439 A	11-06-1998
		WO 9824440 A	11-06-1998
		WO 9824441 A	11-06-1998
		WO 9824443 A	11-06-1998
		WO 9824444 A	11-06-1998
		WO 9824445 A	11-06-1998
		WO 9824446 A	11-06-1998
		WO 9824447 A	11-06-1998
		EP 0906315 A	07-04-1999
		WO 9749710 A	31-12-1997
		HR 970648 A	31-08-1998
		NO 985977 A	22-02-1999
		PL 330235 A	10-05-1999
		US 5929054 A	27-07-1999
		US 5919781 A	06-07-1999
		US 5925627 A	20-07-1999
		AU 8122098 A	19-01-1999
		WO 9900368 A	07-01-1999
WO 9518124 A	06-07-1995	AU 685209 B	15-01-1998
		AU 1322395 A	17-07-1995
		BG 100644 A	31-03-1997
		BR 9408442 A	05-08-1997
		CA 2178219 A	06-07-1995
		CN 1139927 A	08-01-1997
		CZ 9601898 A	11-12-1996
		EP 0737192 A	16-10-1996
		HU 75872 A	28-05-1997
		JP 9507484 T	29-07-1997
		LV 11687 A	20-02-1997
		LV 11687 B	20-10-1997
		NO 962749 A	28-08-1996
		NZ 277839 A	26-01-1998
		PL 315182 A	14-10-1996
		SG 52217 A	28-09-1998
		SK 83996 A	06-11-1996
		US 5612337 A	18-03-1997
		ZA 9410317 A	10-10-1995
		FI 951762 A	13-10-1995
		HR 950222 A	31-08-1997
WO 9718206 A	22-05-1997	AU 7578696 A	05-06-1997
		EP 0861251 A	02-09-1998

THIS PAGE BLANK (USPTO)